

(e) FDA, for good cause, may waive a requirement for the submission of evidence of in vivo bioavailability or bioequivalence if waiver is compatible with the protection of the public health. For full new drug applications, FDA may defer a requirement for the submission of evidence of in vivo bioavailability if deferral is compatible with the protection of the public health.

(f) FDA, for good cause, may require evidence of in vivo bioavailability or bioequivalence for any drug product if the agency determines that any difference between the drug product and a listed drug may affect the bioavailability or bioequivalence of the drug product.

[57 FR 17998, Apr. 28, 1992, as amended at 67 FR 77673, Dec. 19, 2002]

§ 320.23 Basis for measuring in vivo bioavailability or demonstrating bioequivalence.

(a)(1) The in vivo bioavailability of a drug product is measured if the product's rate and extent of absorption, as determined by comparison of measured parameters, e.g., concentration of the active drug ingredient in the blood, urinary excretion rates, or pharmacological effects, do not indicate a significant difference from the reference material's rate and extent of absorption. For drug products that are not intended to be absorbed into the bloodstream, bioavailability may be assessed by measurements intended to reflect the rate and extent to which the active ingredient or active moiety becomes available at the site of action.

(2) Statistical techniques used shall be of sufficient sensitivity to detect differences in rate and extent of absorption that are not attributable to subject variability.

(3) A drug product that differs from the reference material in its rate of absorption, but not in its extent of absorption, may be considered to be bioavailable if the difference in the rate of absorption is intentional, is appropriately reflected in the labeling, is not essential to the attainment of effective body drug concentrations on chronic use, and is considered medically insignificant for the drug product.

(b) Two drug products will be considered bioequivalent drug products if they are pharmaceutical equivalents or pharmaceutical alternatives whose rate and extent of absorption do not show a significant difference when administered at the same molar dose of the active moiety under similar experimental conditions, either single dose or multiple dose. Some pharmaceutical equivalents or pharmaceutical alternatives may be equivalent in the extent of their absorption but not in their rate of absorption and yet may be considered bioequivalent because such differences in the rate of absorption are intentional and are reflected in the labeling, are not essential to the attainment of effective body drug concentrations on chronic use, and are considered medically insignificant for the particular drug product studied.

[57 FR 17999, Apr. 28, 1992, as amended at 67 FR 77673, Dec. 19, 2002]

§ 320.24 Types of evidence to measure bioavailability or establish bioequivalence.

(a) Bioavailability may be measured or bioequivalence may be demonstrated by several in vivo and in vitro methods. FDA may require in vivo or in vitro testing, or both, to measure the bioavailability of a drug product or establish the bioequivalence of specific drug products. Information on bioequivalence requirements for specific products is included in the current edition of FDA's publication "Approved Drug Products with Therapeutic Equivalence Evaluations" and any current supplement to the publication. The selection of the method used to meet an in vivo or in vitro testing requirement depends upon the purpose of the study, the analytical methods available, and the nature of the drug product. Applicants shall conduct bioavailability and bioequivalence testing using the most accurate, sensitive, and reproducible approach available among those set forth in paragraph (b) of this section. The method used must be capable of measuring bioavailability or establishing bioequivalence, as appropriate, for the product being tested.

(b) The following in vivo and in vitro approaches, in descending order of accuracy, sensitivity, and reproducibility, are acceptable for determining the bioavailability or bioequivalence of a drug product.

(1)(i) An in vivo test in humans in which the concentration of the active ingredient or active moiety, and, when appropriate, its active metabolite(s), in whole blood, plasma, serum, or other appropriate biological fluid is measured as a function of time. This approach is particularly applicable to dosage forms intended to deliver the active moiety to the bloodstream for systemic distribution within the body; or

(ii) An in vitro test that has been correlated with and is predictive of human in vivo bioavailability data; or

(2) An in vivo test in humans in which the urinary excretion of the active moiety, and, when appropriate, its active metabolite(s), are measured as a function of time. The intervals at which measurements are taken should ordinarily be as short as possible so that the measure of the rate of elimination is as accurate as possible. Depending on the nature of the drug product, this approach may be applicable to the category of dosage forms described in paragraph (b)(1)(i) of this section. This method is not appropriate where urinary excretion is not a significant mechanism of elimination.

(3) An in vivo test in humans in which an appropriate acute pharmacological effect of the active moiety, and, when appropriate, its active metabolite(s), are measured as a function of time if such effect can be measured with sufficient accuracy, sensitivity, and reproducibility. This approach is applicable to the category of dosage forms described in paragraph (b)(1)(i) of this section only when appropriate methods are not available for measurement of the concentration of the moiety, and, when appropriate, its active metabolite(s), in biological fluids or excretory products but a method is available for the measurement of an appropriate acute pharmacological effect. This approach may be particularly applicable to dosage forms that are not intended to deliver the active moiety

to the bloodstream for systemic distribution.

(4) Well-controlled clinical trials that establish the safety and effectiveness of the drug product, for purposes of measuring bioavailability, or appropriately designed comparative clinical trials, for purposes of demonstrating bioequivalence. This approach is the least accurate, sensitive, and reproducible of the general approaches for measuring bioavailability or demonstrating bioequivalence. For dosage forms intended to deliver the active moiety to the bloodstream for systemic distribution, this approach may be considered acceptable only when analytical methods cannot be developed to permit use of one of the approaches outlined in paragraphs (b)(1)(i) and (b)(2) of this section, when the approaches described in paragraphs (b)(1)(ii), (b)(1)(iii), and (b)(3) of this section are not available. This approach may also be considered sufficiently accurate for measuring bioavailability or demonstrating bioequivalence of dosage forms intended to deliver the active moiety locally, e.g., topical preparations for the skin, eye, and mucous membranes; oral dosage forms not intended to be absorbed, e.g., an antacid or radiopaque medium; and bronchodilators administered by inhalation if the onset and duration of pharmacological activity are defined.

(5) A currently available in vitro test acceptable to FDA (usually a dissolution rate test) that ensures human in vivo bioavailability.

(6) Any other approach deemed adequate by FDA to measure bioavailability or establish bioequivalence.

(c) FDA may, notwithstanding prior requirements for measuring bioavailability or establishing bioequivalence, require in vivo testing in humans of a product at any time if the agency has evidence that the product:

(1) May not produce therapeutic effects comparable to a pharmaceutical equivalent or alternative with which it is intended to be used interchangeably;

(2) May not be bioequivalent to a pharmaceutical equivalent or alternative with which it is intended to be used interchangeably; or

(3) Has greater than anticipated potential toxicity related to pharmacokinetic or other characteristics.

[57 FR 17999, Apr. 28, 1992; 57 FR 29354, July 1, 1992, as amended at 67 FR 77673, Dec. 19, 2002]

§ 320.25 Guidelines for the conduct of an in vivo bioavailability study.

(a) *Guiding principles.* (1) The basic principle in an in vivo bioavailability study is that no unnecessary human research should be done.

(2) An in vivo bioavailability study is generally done in a normal adult population under standardized conditions. In some situations, an in vivo bioavailability study in humans may preferably and more properly be done in suitable patients. Critically ill patients shall not be included in an in vivo bioavailability study unless the attending physician determines that there is a potential benefit to the patient.

(b) *Basic design.* The basic design of an in vivo bioavailability study is determined by the following:

(1) The scientific questions to be answered.

(2) The nature of the reference material and the dosage form to be tested.

(3) The availability of analytical methods.

(4) Benefit-risk considerations in regard to testing in humans.

(c) *Comparison to a reference material.* In vivo bioavailability testing of a drug product shall be in comparison to an appropriate reference material unless some other approach is more appropriate for valid scientific reasons.

(d) *Previously unmarketed active drug ingredients or therapeutic moieties.* (1) An in vivo bioavailability study involving a drug product containing an active drug ingredient or therapeutic moiety that has not been approved for marketing can be used to measure the following pharmacokinetic data:

(i) The bioavailability of the formulation proposed for marketing; and

(ii) The essential pharmacokinetic characteristics of the active drug ingredient or therapeutic moiety, such as the rate of absorption, the extent of absorption, the half-life of the therapeutic moiety in vivo, and the rate of excretion and/or metabolism. Dose proportionality of the active drug ingre-

dient or the therapeutic moiety needs to be established after single-dose administration and in certain instances after multiple-dose administration. This characterization is a necessary part of the investigation of the drug to support drug labeling.

(2) The reference material in such a bioavailability study should be a solution or suspension containing the same quantity of the active drug ingredient or therapeutic moiety as the formulation proposed for marketing.

(3) The reference material should be administered by the same route as the formulation proposed for marketing unless an alternative or additional route is necessary to answer the scientific question under study. For example, in the case of an active drug ingredient or therapeutic moiety that is poorly absorbed after oral administration, it may be necessary to compare the oral dosage form proposed for marketing with the active drug ingredient or therapeutic moiety administered in solution both orally and intravenously.

(e) *New formulations of active drug ingredients or therapeutic moieties approved for marketing.* (1) An in vivo bioavailability study involving a drug product that is a new dosage form, or a new salt or ester of an active drug ingredient or therapeutic moiety that has been approved for marketing can be used to:

(i) Measure the bioavailability of the new formulation, new dosage form, or new salt or ester relative to an appropriate reference material; and

(ii) Define the pharmacokinetic parameters of the new formulation, new dosage form, or new salt or ester to establish dosage recommendation.

(2) The selection of the reference material(s) in such a bioavailability study depends upon the scientific questions to be answered, the data needed to establish comparability to a currently marketed drug product, and the data needed to establish dosage recommendations.

(3) The reference material should be taken from a current batch of a drug product that is the subject of an approved new drug application and that contains the same active drug ingredient or therapeutic moiety, if the new formulation, new dosage form, or new